

# ON THE COHOMOLOGY RING OF ASSOCIATED FLAG BUNDLE

CHEN HE

ABSTRACT. In this expository note, we survey the cohomology rings of flag bundles associated to vector bundles in terms of characteristic classes.

## CONTENTS

|                                                                                     |    |
|-------------------------------------------------------------------------------------|----|
| 1. Introduction                                                                     | 1  |
| 2. Cohomology ring of associated flag bundle                                        | 1  |
| 2.1. Cohomology ring of even dimensional Grassmannians and Leray-Hirsch theorem     | 1  |
| 2.2. The splitting principle of computing cohomology ring of associated flag bundle | 4  |
| 2.3. Applications                                                                   | 8  |
| 3. Final discussion                                                                 | 12 |
| References                                                                          | 13 |

## 1. INTRODUCTION

Let  $V \rightarrow B$  be a complex or real vector bundle, we can take the fibre-wise sets of flags to form associated complex or real flag bundle. Similarly, if  $V \rightarrow B$  is an oriented real vector bundle, then we can take the fibre-wise sets of oriented flags to form oriented flag bundle. In the real or oriented case, we will mainly focus on flags of even dimensional subspaces  $0 \subset L_1 \subset L_2 \cdots \subset L_l \subset \mathbb{R}^{2n}$  or  $\mathbb{R}^{2n+1}$ , with dimensions  $0 < 2k_1 < 2k_2 < \cdots < 2k_l < 2n$  or  $2n+1$ . According to the Leray-Borel description on cohomology of homogeneous spaces, the cohomologies of flag manifolds are generated by Chern classes in the complex case, or by Pontryagin classes and Euler classes in the real case. In this expository note, we follow Bott and Tu's ideas of using splitting principle to give a similar description on the cohomology rings of associated flag bundles.

**Acknowledgement** The author would like to thank his advisors Prof. Victor Guillemin and Prof. Jonathan Weitsman for guidance. The author would also like to thank Jeffrey Carlson for many helpful discussions and references.

## 2. COHOMOLOGY RING OF ASSOCIATED FLAG BUNDLE

### 2.1. Cohomology ring of even dimensional Grassmannians and Leray-Hirsch theorem.

**Theorem 2.1.1** (Leray-Borel description of cohomology of homogeneous spaces, [Le49, Bo53a]). *For compact connected Lie group  $G$  and its connected closed subgroup  $H$  containing the same maximal torus  $T$ , denote*

---

2010 *Mathematics Subject Classification.* Primary 57R22; Secondary 14M15.

*Key words and phrases.* Grassmannian, Flag manifold, Vector bundle, Characteristic classes.

Supported in part by NSF grant DMS 12/11819.

the Weyl groups of  $G, H$  as  $W_G, W_H$ , and the symmetric algebra of the dual Lie algebra  $\mathfrak{t}^*$  as  $\mathfrak{St}^*$ , then the cohomology of homogeneous space  $G/H$  in  $\mathbb{Q}$  coefficients can be given as:

$$H^*(G/H) = (\mathfrak{St}^*)^{W_H} / (\mathfrak{St}^*)_+^{W_G}$$

where  $(\mathfrak{St}^*)^{W_H}$  is the set of  $W_H$  invariants, and  $(\mathfrak{St}^*)_+^{W_G}$  is the set of positive-degree  $W_G$  invariants.

If we apply this general theorem to complex Grassmannian and even dimensional real or oriented Grassmannians, then it is not hard to see that the invariants of Weyl groups are exactly Chern classes, Pontryagin classes and Euler classes of the canonical bundle and complementary bundle of the Grassmannians, with relations from Whitney product formula and square formula of Euler class as top Pontryagin class. More specifically,

**Proposition 2.1.1** (Leray-Borel description of complex Grassmannian, see [BT82]). *For the complex Grassmannian  $G_k(\mathbb{C}^n)$ , let  $c_1, c_2, \dots, c_k$  and  $\bar{c}_1, \bar{c}_2, \dots, \bar{c}_{n-k}$  be the Chern classes of the canonical bundle on  $G_k(\mathbb{C}^n)$  and its complementary bundle respectively, then*

$$H^*(G_k(\mathbb{C}^n), \mathbb{Z}) = \frac{\mathbb{Z}[c_1, c_2, \dots, c_k; \bar{c}_1, \bar{c}_2, \dots, \bar{c}_{n-k}]}{c\bar{c} = 1}$$

where  $c = 1 + c_1 + c_2 + \dots + c_k$ ,  $\bar{c} = 1 + \bar{c}_1 + \bar{c}_2 + \dots + \bar{c}_{n-k}$  are the total Chern classes. Moreover, the Poincaré series are

$$P_{G_k(\mathbb{C}^n)}(t) = \frac{(1-t^2) \cdots (1-t^{2n})}{(1-t^2) \cdots (1-t^{2k})(1-t^2) \cdots (1-t^{2(n-k)})}$$

**Proposition 2.1.2** (Characteristic basis of complex Grassmannian, [Ca78, Ja89]). *The set of monomials  $c_1^{r_1} c_2^{r_2} \cdots c_k^{r_k}$  satisfying the condition  $\sum_{i=1}^k r_i \leq n - k$  forms an additive basis for  $H^*(G_k(\mathbb{C}^n), \mathbb{Z})$ .*

**Example 2.1.1.** The cohomology ring of complex projective space  $\mathbb{C}P^{n-1} = G_1(\mathbb{C}^n)$  can be given as

$$H^*(\mathbb{C}P^{n-1}) = \frac{\mathbb{Z}[c_1; \bar{c}_1, \bar{c}_2, \dots, \bar{c}_{n-1}]}{c\bar{c} = 1} = \mathbb{Z}[c_1]/c_1^n$$

with additive basis  $1, c_1, \dots, c_1^{n-1}$  and Poincaré series  $(1-t^{2n})/(1-t^2)$ .

**Proposition 2.1.3** (Leray-Borel description of even dimensional real Grassmannians, see [He]). *For the even dimensional real Grassmannians  $G_{2k}(\mathbb{R}^{2n})$ ,  $G_{2k}(\mathbb{R}^{2n+1})$  and  $G_{2k+1}(\mathbb{R}^{2n+1})$ , let  $p_1, p_2, \dots, p_k$  and  $\bar{p}_1, \bar{p}_2, \dots, \bar{p}_{n-k}$  be the Pontryagin classes of canonical bundles and complementary bundles over these Grassmannians, their cohomologies in  $\mathbb{Q}$  coefficients are the same:*

$$H^*(G_{2k}(\mathbb{R}^{2n}), \mathbb{Q}) \cong H^*(G_{2k}(\mathbb{R}^{2n+1}), \mathbb{Q}) \cong H^*(G_{2k+1}(\mathbb{R}^{2n+1}), \mathbb{Q}) = \frac{\mathbb{Q}[p_1, p_2, \dots, p_k; \bar{p}_1, \bar{p}_2, \dots, \bar{p}_{n-k}]}{p\bar{p} = 1}$$

**Proposition 2.1.4** (Characteristic basis of even dimensional real Grassmannians, [He]). *The set of monomials  $(p_1)^{r_1} (p_2)^{r_2} \cdots (p_k)^{r_k}$  satisfying the condition  $\sum_{i=1}^k r_i \leq n - k$  forms an additive basis for  $H^*(G_{2k}(\mathbb{R}^{2n}), \mathbb{Q}) \cong H^*(G_{2k}(\mathbb{R}^{2n+1}), \mathbb{Q}) \cong H^*(G_{2k+1}(\mathbb{R}^{2n+1}), \mathbb{Q})$ . Moreover, the Poincaré series are all equal to  $P_{G_k(\mathbb{C}^n)}(t^2)$ .*

*Remark 2.1.1.* To the best knowledge of the author, the Poincaré series of real Grassmannians were first explicitly given by Casian and Kodama [CK].

**Example 2.1.2.** The cohomology ring of  $\mathbb{R}P^{2n} = G_1(\mathbb{R}^{2n+1})$  can be given as

$$H^*(\mathbb{R}P^{2n}, \mathbb{Q}) = \frac{\mathbb{Q}[\bar{p}_1, \bar{p}_2, \dots, \bar{p}_n]}{\bar{p} = 1} = \mathbb{Q}$$

**Example 2.1.3.** The cohomology rings of real Grassmannians  $G_2(\mathbb{R}^{2n})$ ,  $G_2(\mathbb{R}^{2n+1})$  and  $G_3(\mathbb{R}^{2n+1})$  can be given as

$$\begin{aligned} H^*(G_2(\mathbb{R}^{2n}), \mathbb{Q}) &\cong H^*(G_2(\mathbb{R}^{2n+1}), \mathbb{Q}) \cong H^*(G_3(\mathbb{R}^{2n+1}), \mathbb{Q}) \\ &= \frac{\mathbb{Q}[p_1; \bar{p}_1, \bar{p}_2, \dots, \bar{p}_{n-1}]}{p\bar{p} = 1} = \mathbb{Q}[p_1]/p_1^n \end{aligned}$$

with additive basis  $1, p_1, \dots, p_1^{n-1}$  and Poincaré series  $(1 - t^{4n})/(1 - t^4)$ .

**Proposition 2.1.5** (Leray-Borel description of even dimensional oriented Grassmannians, see [He]). *For the even dimensional oriented Grassmannians  $\tilde{G}_{2k}(\mathbb{R}^{2n})$ ,  $\tilde{G}_{2k}(\mathbb{R}^{2n+1})$  and  $\tilde{G}_{2k+1}(\mathbb{R}^{2n+1})$ , besides the Pontryagin classes of canonical bundles and complementary bundles, let  $e$  be the Euler class of canonical bundles over  $\tilde{G}_{2k}(\mathbb{R}^{2n})$ ,  $\tilde{G}_{2k}(\mathbb{R}^{2n+1})$ , and let  $\bar{e}$  be the Euler class of complementary bundles over  $\tilde{G}_{2k}(\mathbb{R}^{2n})$ ,  $\tilde{G}_{2k+1}(\mathbb{R}^{2n+1})$ , their cohomologies in  $\mathbb{Q}$  coefficients are given as:*

$$\begin{aligned} H^*(\tilde{G}_{2k}(\mathbb{R}^{2n}), \mathbb{Q}) &= \frac{\mathbb{Q}[p_1, p_2, \dots, p_k; \bar{p}_1, \bar{p}_2, \dots, \bar{p}_{n-k}; e, \bar{e}]}{p\bar{p} = 1, e^2 = p_k, \bar{e}^2 = \bar{p}_{n-k}, e\bar{e} = 0} \\ H^*(\tilde{G}_{2k}(\mathbb{R}^{2n+1}), \mathbb{Q}) &= \frac{\mathbb{Q}[p_1, p_2, \dots, p_k; \bar{p}_1, \bar{p}_2, \dots, \bar{p}_{n-k}; e]}{p\bar{p} = 1, e^2 = p_k} \\ H^*(\tilde{G}_{2k+1}(\mathbb{R}^{2n+1}), \mathbb{Q}) &= \frac{\mathbb{Q}[p_1, p_2, \dots, p_k; \bar{p}_1, \bar{p}_2, \dots, \bar{p}_{n-k}; \bar{e}]}{p\bar{p} = 1, \bar{e}^2 = \bar{p}_{n-k}} \end{aligned}$$

**Proposition 2.1.6** (Characteristic basis of even dimensional oriented Grassmannians, [He]). *The sets of monomials  $\{p_1^{r_1} p_2^{r_2} \dots p_k^{r_k}, e \cdot p_1^{r_1} p_2^{r_2} \dots p_k^{r_k}\}$  and  $\{p_1^{r_1} p_2^{r_2} \dots p_k^{r_k}, \bar{e} \cdot p_1^{r_1} p_2^{r_2} \dots p_k^{r_k}\}$  satisfying the condition  $\sum_{i=1}^k r_i \leq n - k$  form additive basis for  $H^*(\tilde{G}_{2k}(\mathbb{R}^{2n+1}), \mathbb{Q})$ ,  $H^*(\tilde{G}_{2k+1}(\mathbb{R}^{2n+1}), \mathbb{Q})$  respectively. The set of monomials  $\{p_1^{r_1} p_2^{r_2} \dots p_k^{r_k} \mid \sum_{i=1}^k r_i \leq n - k\} \cup \{e \cdot p_1^{r_1} p_2^{r_2} \dots p_{n-k-1}^{r_{n-k-1}} \mid \sum_{i=1}^{n-k-1} r_i \leq k\} \cup \{\bar{e} \cdot p_1^{r_1} p_2^{r_2} \dots p_{k-1}^{r_{k-1}} \mid \sum_{i=1}^{k-1} r_i \leq n - k\}$  forms additive basis for  $H^*(\tilde{G}_{2k}(\mathbb{R}^{2n}), \mathbb{Q})$ . Moreover, the Poincaré series are:*

$$\begin{aligned} P_{\tilde{G}_{2k}(\mathbb{R}^{2n})}(t) &= P_{G_k(\mathbb{C}^n)}(t^2) + t^{2k} P_{G_k(\mathbb{C}^{n-1})}(t^2) + t^{2n-2k} P_{G_{k-1}(\mathbb{C}^{n-1})}(t^2) \\ P_{\tilde{G}_{2k}(\mathbb{R}^{2n+1})}(t) &= (1 + t^{2k}) P_{G_k(\mathbb{C}^n)}(t^2) \\ P_{\tilde{G}_{2k+1}(\mathbb{R}^{2n+1})}(t) &= (1 + t^{2n-2k}) P_{G_k(\mathbb{C}^n)}(t^2) \end{aligned}$$

*Remark 2.1.2.* The Poincaré series of even dimensional oriented Grassmannians were already computed by H. Cartan [Car50].

**Example 2.1.4.** The cohomology ring of  $S^{2n} = \tilde{G}_1(\mathbb{R}^{2n+1})$  can be given as

$$H^*(S^{2n}, \mathbb{Q}) = \frac{\mathbb{Q}[\bar{p}_1, \bar{p}_2, \dots, \bar{p}_n; \bar{e}]}{\bar{p} = 1, \bar{e}^2 = \bar{p}_n} = \mathbb{Q}[\bar{e}]/\bar{e}^2$$

where  $\bar{e} \in H^{2n}$  is the Euler class of the tangent bundle over  $S^{2n}$  and is also an orientation form of  $S^{2n}$ , with additive basis  $1, \bar{e}$  and Poincaré series  $1 + t^{2n}$ .

**Example 2.1.5.** The cohomology ring of oriented Grassmannian  $\tilde{G}_2(\mathbb{R}^{2n})$  can be given as

$$\begin{aligned} H^*(\tilde{G}_2(\mathbb{R}^{2n}), \mathbb{Q}) &= \frac{\mathbb{Q}[p_1; \bar{p}_1, \bar{p}_2, \dots, \bar{p}_{n-1}; e, \bar{e}]}{p\bar{p} = 1, e^2 = p_1, \bar{e}^2 = \bar{p}_{n-1}, e\bar{e} = 0} \\ &= \frac{\mathbb{Q}[e, \bar{e}]}{e^{2n-1} = 0, e\bar{e} = 0, \bar{e}^2 = (-1)^{n-1} e^{2n-2}} \end{aligned}$$

where  $e \in H^2, \bar{e} \in H^{2n-2}$ , with additive basis  $1, e, \dots, e^{2n-2}, \bar{e}$  and Poincaré series  $t^{2n-2} + (1 - t^{4n-2})/(1 - t^2) = (1 + t^{2n-2})(1 - t^{2n})/(1 - t^2)$ .

**Example 2.1.6.** The cohomology ring of oriented Grassmannian  $\tilde{G}_2(\mathbb{R}^{2n+1})$  can be given as

$$H^*(\tilde{G}_2(\mathbb{R}^{2n+1}), \mathbb{Q}) = \frac{\mathbb{Q}[p_1; \bar{p}_1, \bar{p}_2, \dots, \bar{p}_{n-1}; e]}{p\bar{p} = 1, e^2 = p_1} = \mathbb{Q}[e]/e^{2n}$$

where  $e \in H^2$ , with additive basis  $1, e, \dots, e^{2n-1}$  and Poincaré series  $(1 - t^{4n})/(1 - t^2)$ .

**Example 2.1.7.** The cohomology ring of oriented Grassmannian  $\tilde{G}_3(\mathbb{R}^{2n+1})$  can be given as

$$\begin{aligned} H^*(\tilde{G}_3(\mathbb{R}^{2n+1}), \mathbb{Q}) &= \frac{\mathbb{Q}[p_1; \bar{p}_1, \bar{p}_2, \dots, \bar{p}_{n-1}; \bar{e}]}{p\bar{p} = 1, \bar{e}^2 = \bar{p}_{n-1}} \\ &= \frac{\mathbb{Q}[p_1, \bar{e}]}{p_1^n = 0, \bar{e}^2 = (-1)^{n-1} p_1^{n-1}} \end{aligned}$$

where  $\bar{e} \in H^{2n-2}$ , with additive basis  $1, p_1, \dots, p_1^{n-1}; \bar{e}, \bar{e}p_1, \dots, \bar{e}p_1^{n-1}$  and Poincaré series  $(1 + t^{2n-2})(1 - t^{4n})/(1 - t^4)$ .

**Theorem 2.1.2** (Leray-Hirsch isomorphism theorem, see [Ha02]). *Let  $F \xrightarrow{\iota} E \xrightarrow{\pi} B$  be a fibre bundle, such that for some commutative coefficient ring  $R$ :*

- (a)  $H_n(F; R)$  is a finitely generated free  $R$ -module for each  $n$ .
- (b) There exist classes  $c_j \in H^{k_j}(E; R)$  whose restrictions  $\iota^*(c_j)$  form a basis for  $H(F; R)$  in each fiber  $F$ .

Then  $\pi^* : H^*(B; R) \rightarrow H^*(E; R)$  is injective, and the map

$$\begin{aligned} \Phi : H(B; R) \otimes_R H_i(F; R) &\longrightarrow H^*(E; R) \\ \sum_{ij} b_i \otimes \iota^*(c_j) &\longmapsto \sum_{ij} \pi^*(b_i) \smile c_j \end{aligned}$$

is a  $H^*(B; R)$ -module isomorphism and gives  $H^*(E; R)$  a free  $H^*(B; R)$ -module structure with additive  $H^*(B; R)$ -basis  $c_j$ . Moreover, the ring structure of  $H^*(E; R)$  can be specified for certain  $b_{ij}^k \in H(B; R)$  as

$$c_i \smile c_j = \sum_k \pi^*(b_{ij}^k) \smile c_k$$

*Remark 2.1.3.* The Poincaré series of  $E$  are given as  $P_E(t) = P_B(t) \cdot P_F(t)$ .

*Remark 2.1.4.* Under the assumptions of Leray-Hirsch theorem, since  $\pi^* : H^*(B; R) \rightarrow H^*(E; R)$  is a monomorphism, we will often identify every class  $b \in H(B; R)$  with its image  $\pi^*(b) \in H(E; R)$ .

**2.2. The splitting principle of computing cohomology ring of associated flag bundle.** Let  $V \rightarrow B$  be a vector bundle. Without loss of generality, we will compute cohomology ring for associated (even-rank) Grassmannian bundle, and for associated complete (even-rank) flag bundle whose fibre over  $b \in B$  is the set of flags  $0 \subset L_1 \subset L_2 \subset \dots \subset L_{n-1} \subset V_b$  with real dimensions  $0 < 2 < 4 < \dots < 2n - 2 < 2n$  or  $2n + 1$ . We will denote this associated complete even-rank flag bundle as  $Fl(V)$ .

For a rank- $n$  complex or real vector bundle  $\pi_1 : V \rightarrow B$ , let  $\pi_2 : G_k(V) \rightarrow B$  be the associated Grassmannian bundle of  $k$ -dimensional subspaces of  $V$ , then we can pull back the vector bundle  $V$  to be over  $G_k(V)$ , and get a natural splitting of this pull-back bundle as a direct sum of the canonical bundle of  $k$ -dimensional subspaces and the complementary bundle of  $(n - k)$ -dimensional subspaces. The canonical bundle and complementary bundle will be denoted as  $V', V''$  respectively. For any point  $b \in B$  and the vector space  $V_b = \pi_1^{-1}(b)$ , the Grassmannian  $G_k(V_b)$  of  $k$ -dimensional subspaces in  $V_b$  gives a fibre of the Grassmannian bundle  $G_k(V_b) \xrightarrow{\iota} G_k(V) \xrightarrow{\pi_2} B$ . Let's denote  $\gamma', \gamma''$  as the canonical bundle and complementary bundle over  $G_k(V_b)$ . By the universality of canonical bundle and complementary bundle, the inclusion  $\iota$  pulls back  $V', V''$  to be  $\gamma', \gamma''$  over  $G_k(V_b)$ . Therefore, we get a commutative diagram:

$$\begin{array}{ccccc} V_b = \gamma' \oplus \gamma'' & \longrightarrow & \pi_2^* V = V' \oplus V'' & \longrightarrow & V \\ \downarrow & & \downarrow & & \downarrow \pi_1 \\ G_k(V_b) & \xrightarrow{\iota} & G_k(V) & \xrightarrow{\pi_2} & B \end{array}$$

By the naturality of total Chern or Pontryagin classes, we get:

$$\begin{aligned} \pi_2^*(c(V)) &= c(\pi_2^*V) & \iota^*(c(V')) &= c(\gamma') & \iota^*(c(V'')) &= c(\gamma'') \\ \text{or } \pi_2^*(p(V)) &= p(\pi_2^*V) & \iota^*(p(V')) &= p(\gamma') & \iota^*(p(V'')) &= p(\gamma'') \end{aligned}$$

By the Whitney product formula of total Chern or Pontryagin classes, we get:

$$\begin{aligned} c(\pi_2^*V) &= c(V')c(V'') & c(V_b) &= c(\gamma')c(\gamma'') \\ \text{or } p(\pi_2^*V) &= p(V')p(V'') & p(V_b) &= p(\gamma')p(\gamma'') \end{aligned}$$

Similarly, we get the same commutative diagram for the associated oriented Grassmannian bundles of an oriented vector bundle and relations of Euler classes using naturality and Whitney product formula.

**Theorem 2.2.1** (Leray-Borel description of cohomology of associated (even-dim) Grassmannian bundle). *Consider associated Grassmannian bundles of a vector bundle  $V \rightarrow B$ ,*

(1) *If  $V^n$  is a rank- $n$  complex vector bundle, then we get the cohomology ring*

$$H^*(G_k(V^n), \mathbb{Z}) \cong \frac{H^*(B, \mathbb{Z})[c_1, c_2, \dots, c_k; \bar{c}_1, \bar{c}_2, \dots, \bar{c}_{n-k}]}{c\bar{c} = c(V^n)}$$

*with Poincaré series  $P_{G_k(V^n)}(t) = P_B(t)P_{G_k(\mathbb{C}^n)}(t)$ .*

(2) *If  $V$  is a real vector bundle of rank  $2n$  or  $2n+1$ , denoted as  $V^{2n}$  or  $V^{2n+1}$  respectively, then we get the cohomology rings*

$$\begin{aligned} H^*(G_{2k}(V^{2n}), \mathbb{Q}) &\cong \frac{H^*(B, \mathbb{Q})[p_1, p_2, \dots, p_k; \bar{p}_1, \bar{p}_2, \dots, \bar{p}_{n-k}]}{p\bar{p} = p(V^{2n})} \\ H^*(G_{2k}(V^{2n+1}), \mathbb{Q}) &\cong H^*(G_{2k+1}(V^{2n+1}), \mathbb{Q}) \cong \frac{H^*(B, \mathbb{Q})[p_1, p_2, \dots, p_k; \bar{p}_1, \bar{p}_2, \dots, \bar{p}_{n-k}]}{p\bar{p} = p(V^{2n+1})} \end{aligned}$$

*with Poincaré series  $P_{G_{2k}(V^{2n})}(t) = P_B(t)P_{G_{2k}(\mathbb{R}^{2n})}(t)$ , or  $P_{G_{2k}(V^{2n+1})}(t) = P_{G_{2k+1}(V^{2n+1})}(t) = P_B(t)P_{G_{2k}(\mathbb{R}^{2n+1})}(t)$ .*

(3) *If  $V$  is an oriented vector bundle of rank  $2n$  or  $2n+1$ , denoted as  $V^{2n}$  or  $V^{2n+1}$  respectively, then we get the cohomology rings*

$$\begin{aligned} H^*(\tilde{G}_{2k}(V^{2n}), \mathbb{Q}) &\cong \frac{H^*(B, \mathbb{Q})[p_1, p_2, \dots, p_k; \bar{p}_1, \bar{p}_2, \dots, \bar{p}_{n-k}; e, \bar{e}]}{p\bar{p} = p(V^{2n}), e^2 = p_k, \bar{e}^2 = \bar{p}_{n-k}, e\bar{e} = e(V^{2n})} \\ H^*(\tilde{G}_{2k}(V^{2n+1}), \mathbb{Q}) &\cong \frac{H^*(B, \mathbb{Q})[p_1, p_2, \dots, p_k; \bar{p}_1, \bar{p}_2, \dots, \bar{p}_{n-k}; e]}{p\bar{p} = p(V^{2n+1}), e^2 = p_k} \\ H^*(\tilde{G}_{2k+1}(V^{2n+1}), \mathbb{Q}) &\cong \frac{H^*(B, \mathbb{Q})[p_1, p_2, \dots, p_k; \bar{p}_1, \bar{p}_2, \dots, \bar{p}_{n-k}; \bar{e}]}{p\bar{p} = p(V^{2n+1}), \bar{e}^2 = \bar{p}_{n-k}} \end{aligned}$$

*with Poincaré series  $P_{\tilde{G}_{2k}(V^{2n})}(t) = P_B(t)P_{\tilde{G}_{2k}(\mathbb{R}^{2n})}(t)$ , or  $P_{\tilde{G}_{2k}(V^{2n+1})}(t) = P_B(t)P_{\tilde{G}_{2k}(\mathbb{R}^{2n+1})}(t)$  and  $P_{\tilde{G}_{2k+1}(V^{2n+1})}(t) = P_B(t)P_{\tilde{G}_{2k+1}(\mathbb{R}^{2n+1})}(t)$ .*

First of all, we observe that the associated Grassmannian bundles satisfy the assumptions of the Leray-Hirsch Theorem 2.1.2. The condition (a) on the freeness of cohomology of a Grassmannian fibre is guaranteed by the Leray-Borel description 2.1.1 of homogeneous spaces in terms of even-degree characteristic classes. The condition (b) on the existence of preimages of additive basis in cohomology of a Grassmannian fibre is guaranteed by the characteristic basis and naturality of characteristic classes. Moreover, we can specify the ring structure by understanding the multiplications among additive basis.

Secondly, in the real or oriented case, since  $G_{2k+1}(V^{2n+1}) \cong G_{2n-2k}(V^{2n+1})$  and  $\tilde{G}_{2k+1}(V^{2n+1}) \cong \tilde{G}_{2n-2k}(V^{2n+1})$ , we only need to consider associated Grassmannian bundles  $G_{2k}(V^{2n})$ ,  $G_{2k}(V^{2n+1})$  or  $\tilde{G}_{2k}(V^{2n})$ ,  $\tilde{G}_{2k}(V^{2n+1})$ .

2.2.1. *Proof of Theorem 2.2.1.* We will follow Bott&Tu's ideas of using splitting principle to give a three-step proof.

Step 1: Theorem 2.2.1 is true for  $k = 0$  or  $1$ .

**Proposition 2.2.1.** *For a vector bundle  $V \rightarrow B$ ,*

(1) *If  $V^n$  is a rank- $n$  complex vector bundle, then we get the cohomology ring for the associated projectivization bundle*

$$\begin{aligned} H^*(\mathbb{P}(V^n), \mathbb{Z}) &\cong \frac{H^*(B, \mathbb{Z})[c_1; \bar{c}_1, \bar{c}_2, \dots, \bar{c}_{n-1}]}{c\bar{c} = c(V^n)} \\ &\cong \frac{H^*(B, \mathbb{Z})[c_1]}{c_1^n - c_1(V)c_1^{n-1} + c_2(V)c_1^{n-2} \mp \dots + (-1)^n c_n(V) = 0} \end{aligned}$$

(2) *If  $V$  is a real vector bundle of rank  $2n$  or  $2n+1$ , denoted as  $V^{2n}$  or  $V^{2n+1}$  respectively, then we get the cohomology rings for the associated rank-2 Grassmannian bundles*

$$\begin{aligned} H^*(G_2(V^{2n}), \mathbb{Q}) &\cong \frac{H^*(B, \mathbb{Q})[p_1; \bar{p}_1, \bar{p}_2, \dots, \bar{p}_{n-1}]}{p\bar{p} = p(V^{2n})} \\ H^*(G_2(V^{2n+1}), \mathbb{Q}) &\cong \frac{H^*(B, \mathbb{Q})[p_1; \bar{p}_1, \bar{p}_2, \dots, \bar{p}_{n-1}]}{p\bar{p} = p(V^{2n+1})} \end{aligned}$$

*both of which have the same simplified expression:*

$$\frac{H^*(B, \mathbb{Q})[p_1]}{p_1^n - p_1(V)p_1^{n-1} + p_2(V)p_1^{n-2} \mp \dots + (-1)^n p_n(V) = 0}$$

(3) *If  $V$  is an oriented vector bundle of rank  $2n$  or  $2n+1$ , denoted as  $V^{2n}$  or  $V^{2n+1}$  respectively, then we get the cohomology rings for the associated sphere or rank-2 oriented Grassmannian bundles*

$$\begin{aligned} H^*(\tilde{G}_2(V^{2n}), \mathbb{Q}) &\cong \frac{H^*(B, \mathbb{Q})[p_1; \bar{p}_1, \bar{p}_2, \dots, \bar{p}_{n-1}; e, \bar{e}]}{p\bar{p} = p(V^{2n}), e^2 = p_1, \bar{e}^2 = \bar{p}_{n-1}, e\bar{e} = e(V^{2n})} \\ H^*(S(V^{2n+1}), \mathbb{Q}) &\cong \frac{H^*(B, \mathbb{Q})[\bar{p}_1, \bar{p}_2, \dots, \bar{p}_n; \bar{e}]}{\bar{p} = p(V^{2n+1}), \bar{e}^2 = \bar{p}_n} \cong \frac{H^*(B, \mathbb{Q})[\bar{e}]}{\bar{e}^2 \cong p_n(V^{2n+1})} \\ H^*(\tilde{G}_2(V^{2n+1}), \mathbb{Q}) &\cong \frac{H^*(B, \mathbb{Q})[p_1; \bar{p}_1, \bar{p}_2, \dots, \bar{p}_{n-1}; e]}{p\bar{p} = p(V^{2n+1}), e^2 = p_1} \\ &\cong \frac{H^*(B, \mathbb{Q})[e]}{e^{2n} - p_1(V)e^{2n-2} + p_2(V)e^{2n-4} \mp \dots + (-1)^n p_n(V) = 0} \end{aligned}$$

*Proof.* This proposition follows directly from the explicitly worked out Examples 2.1.1-2.1.7, the naturality of characteristic classes and Leray-Hirsch theorem.  $\square$

Step 2: Similar Theorem is true for associated complete even-rank flag bundles.

**Proposition 2.2.2.** *For a vector bundle  $V \rightarrow B$ ,*

(1) *If  $V^n$  is a rank- $n$  complex vector bundle, then we get the cohomology ring for the associated complete complex flag bundle*

$$H^*(Fl(V^n), \mathbb{Z}) \cong \frac{H^*(B, \mathbb{Z})[x_1, x_2, \dots, x_n]}{(1+x_1)(1+x_2)\dots(1+x_n) = c(V^n)}$$

*where  $x_i \in H^2$  is the first Chern class of the  $i$ -th canonical complex line bundle over  $Fl(V)$ . Moreover, the Poincaré series are*

$$P_{Fl(V)}(t) = P_B(t) \prod_{i=1}^n P_{\mathbb{C}P^{i-1}}(t) = P_B(t) \prod_{i=1}^n \frac{1-t^{2i}}{1-t^2}$$

- (2) If  $V$  is a real vector bundle of rank  $2n$  or  $2n + 1$ , denoted as  $V^{2n}$  or  $V^{2n+1}$  respectively, then we get the cohomology rings for the associated complete even-rank flag bundles

$$H^*(Fl(V^{2n}), \mathbb{Q}) \cong \frac{H^*(B, \mathbb{Q})[u_1, u_2, \dots, u_n]}{(1 + u_1)(1 + u_2) \cdots (1 + u_n) = p(V^{2n})}$$

$$H^*(Fl(V^{2n+1}), \mathbb{Q}) \cong \frac{H^*(B, \mathbb{Q})[u_1, u_2, \dots, u_n]}{(1 + u_1)(1 + u_2) \cdots (1 + u_n) = p(V^{2n+1})}$$

where  $u_i \in H^4$  is the first Pontryagin class of the  $i$ -th canonical rank-2 vector bundle over  $Fl(V)$ . Moreover, the Poincaré series are

$$P_{Fl(V)}(t) = P_B(t) \prod_{i=1}^n P_{G_2(\mathbb{R}^{2i})}(t) = P_B(t) \prod_{i=1}^n \frac{1 - t^{4i}}{1 - t^4}$$

- (3) If  $V$  is an oriented vector bundle of rank  $2n$  or  $2n + 1$ , denoted as  $V^{2n}$  or  $V^{2n+1}$  respectively, then we get the cohomology rings for the associated complete even-rank oriented flag bundles

$$H^*(\tilde{Fl}(V^{2n}), \mathbb{Q}) \cong \frac{H^*(B, \mathbb{Q})[u_1, u_2, \dots, u_n; e_1, e_2, \dots, e_n]}{(1 + u_1)(1 + u_2) \cdots (1 + u_n) = p(V^{2n}), e_i^2 = u_i, e_1 e_2 \cdots e_n = e(V^{2n})}$$

$$\cong \frac{H^*(B, \mathbb{Q})[e_1, e_2, \dots, e_n]}{(1 + e_1^2)(1 + e_2^2) \cdots (1 + e_n^2) = p(V^{2n}), e_1 e_2 \cdots e_n = e(V^{2n})}$$

$$H^*(\tilde{Fl}(V^{2n+1}), \mathbb{Q}) \cong \frac{H^*(B, \mathbb{Q})[u_1, u_2, \dots, u_n; e_1, e_2, \dots, e_n]}{(1 + u_1)(1 + u_2) \cdots (1 + u_n) = p(V^{2n}), e_i^2 = u_i}$$

$$\cong \frac{H^*(B, \mathbb{Q})[e_1, e_2, \dots, e_n]}{(1 + e_1^2)(1 + e_2^2) \cdots (1 + e_n^2) = p(V^{2n+1})}$$

where  $e_i \in H^2, u_i \in H^4$  are respectively the Euler class and the first Pontryagin class of the  $i$ -th canonical rank-2 vector bundle over  $Fl(V)$ . Moreover, the Poincaré series are

$$P_{\tilde{Fl}(V^{2n})}(t) = P_B(t) \prod_{i=1}^n P_{\tilde{G}_2(\mathbb{R}^{2i})}(t) = P_B(t) \prod_{i=1}^n \frac{(1 + t^{2i-2})(1 - t^{2i})}{1 - t^2}$$

$$P_{\tilde{Fl}(V^{2n+1})}(t) = P_B(t) \prod_{i=1}^n P_{\tilde{G}_2(\mathbb{R}^{2i+1})}(t) = P_B(t) \prod_{i=1}^n \frac{1 - t^{4i}}{1 - t^2}$$

*Proof.* Note the associated complete flag bundles are obtained through a series of projectivizations in the complex case or rank-2 Grassmannianizations in the real or oriented case. The current proposition then follows from the previous Proposition 2.2.1. The computation of Poincaré series follows from Remark 2.1.3 and the Poincaré series in Examples 2.1.1-2.1.7.  $\square$

Step 3: Theorem 2.2.1 is true for any  $k \leq n$ : for any associated (even-rank) Grassmannian bundle, both of its canonical vector bundle and complementary vector bundle have associated complete (even-rank) flag bundles, then Step 3 follows from Step 2.

*Remark 2.2.1.* In Bott and Tu's book [BT82], Steps 1&2 are their Proposition 21.16 and its following paragraphs. Step 3 is their Proposition 23.2.

*Remark 2.2.2.* The same ideas also apply to the computation of cohomology ring of associated partial (even-rank) flag bundle.

*Remark 2.2.3.* In  $\mathbb{Z}/2$  coefficients, there is also Leray-Borel descriptions on the cohomology of real Grassmannian in [Bo53b]. We can consider associated real partial flag bundle with general flags not necessarily of even rank. The cohomology ring of associated real partial flag bundle is the ring extension of cohomology of the base by the Stiefel-Whitney classes with relations from the Whitney product formula. For associated complex partial

flag bundle, we then replace Chern classes  $c_i$  by even-degree Stiefel-Whitney classes  $w_{2i}$  and ignore the odd-degree Stiefel-Whitney classes. However, the above calculation of cohomology ring does not seem to apply to associated oriented partial flag bundle, because the cohomology of oriented Grassmannian in  $\mathbb{Z}/2$  coefficients is still unknown, to the best knowledge of the author.

**2.3. Applications.** We give some applications of the Theorem 2.2.1, Propositions 2.2.1, 2.2.2 and Remark 2.2.2.

**2.3.1. Complete flag manifold.** Take the base  $B$  to be a point, and the vector bundle  $V$  to be a complex or real vector space. From Proposition 2.2.2:

- (1) If  $V = \mathbb{C}^n$ , then we get the cohomology ring for the complete complex flag manifold

$$H^*(Fl(\mathbb{C}^n), \mathbb{Z}) = \frac{\mathbb{Z}[x_1, x_2, \dots, x_n]}{(1+x_1)(1+x_2)\cdots(1+x_n)=1}$$

where  $x_i \in H^2$  is the first Chern class of the  $i$ -th canonical complex line bundle over  $Fl(\mathbb{C}^n)$ . Moreover, the Poincaré series are

$$P_{Fl(\mathbb{C}^n)}(t) = \prod_{i=1}^n \frac{1-t^{2i}}{1-t^2}$$

- (2) If  $V = \mathbb{R}^{2n}$  or  $\mathbb{R}^{2n+1}$ , then we get the cohomology rings for the complete even-rank flag manifolds

$$\begin{aligned} H^*(Fl(\mathbb{R}^{2n}), \mathbb{Q}) &= \frac{\mathbb{Q}[u_1, u_2, \dots, u_n]}{(1+u_1)(1+u_2)\cdots(1+u_n)=1} \\ H^*(Fl(\mathbb{R}^{2n+1}), \mathbb{Q}) &= \frac{\mathbb{Q}[u_1, u_2, \dots, u_n]}{(1+u_1)(1+u_2)\cdots(1+u_n)=1} \end{aligned}$$

where  $u_i \in H^4$  is the first Pontryagin class of the  $i$ -th canonical rank-2 vector bundle over  $Fl(V)$ . Moreover, the Poincaré series are

$$P_{Fl(\mathbb{R}^{2n})}(t) = P_{Fl(\mathbb{R}^{2n+1})}(t) = \prod_{i=1}^n \frac{1-t^{4i}}{1-t^4}$$

- (3) If  $V = \mathbb{R}^{2n}$  or  $\mathbb{R}^{2n+1}$  with the natural orientation, then we get the cohomology rings for the complete even-rank oriented flag manifolds

$$\begin{aligned} H^*(\tilde{Fl}(\mathbb{R}^{2n}), \mathbb{Q}) &= \frac{\mathbb{Q}[u_1, u_2, \dots, u_n; e_1, e_2, \dots, e_n]}{(1+u_1)(1+u_2)\cdots(1+u_n)=1, e_i^2=u_i, e_1e_2\cdots e_n=0} \\ &= \frac{\mathbb{Q}[e_1, e_2, \dots, e_n]}{(1+e_1^2)(1+e_2^2)\cdots(1+e_n^2)=1, e_1e_2\cdots e_n=0} \\ H^*(\tilde{Fl}(\mathbb{R}^{2n+1}), \mathbb{Q}) &= \frac{\mathbb{Q}[u_1, u_2, \dots, u_n; e_1, e_2, \dots, e_n]}{(1+u_1)(1+u_2)\cdots(1+u_n)=1, e_i^2=u_i} \\ &= \frac{\mathbb{Q}[e_1, e_2, \dots, e_n]}{(1+e_1^2)(1+e_2^2)\cdots(1+e_n^2)=1} \end{aligned}$$

where  $e_i \in H^2$  is the Euler class of the  $i$ -th canonical rank-2 bundle over  $\tilde{Fl}(V)$ . Moreover, the Poincaré series are

$$P_{\tilde{Fl}(\mathbb{R}^{2n})}(t) = \prod_{i=1}^n \frac{(1+t^{2i-2})(1-t^{2i})}{1-t^2} \quad P_{\tilde{Fl}(\mathbb{R}^{2n+1})}(t) = \prod_{i=1}^n \frac{1-t^{4i}}{1-t^2}$$

2.3.2. *The odd dimensional Grassmannians  $G_{2k+1}(\mathbb{R}^{2n+2})$  and  $\tilde{G}_{2k+1}(\mathbb{R}^{2n+2})$ .* Consider the fibre bundle

$$\frac{O(2k+1)}{O(2)^{2k} \times O(1)} \times \frac{O(2n-2k+1)}{O(2)^{2n-2k} \times O(1)} \hookrightarrow \frac{O(2n+2)}{O(2)^{2k} \times O(1) \times O(2)^{2n-2k} \times O(1)} \rightarrow \frac{O(2n+2)}{O(2k+1) \times O(2n-2k+1)}$$

where the fibre is the product of complete even-rank real flag manifolds  $Fl(\mathbb{R}^{2k+1}) \times Fl(\mathbb{R}^{2n-2k+1})$  and the base is the odd dimensional Grassmannian  $G_{2k+1}(\mathbb{R}^{2n+2})$ . Applying the Proposition 2.2.2 to this complete even-rank flag bundle, we get:

$$H^*\left(\frac{O(2n+2)}{O(2)^{2k} \times O(1) \times O(2)^{2n-2k} \times O(1)}, \mathbb{Q}\right) \cong \frac{H^*(G_{2k+1}(\mathbb{R}^{2n+2}), \mathbb{Q})[u_1, \dots, u_k; u_{k+1}, \dots, u_n]}{\prod_{i=1}^k (1+u_i) = p(\gamma), \prod_{j=1}^{n-k} (1+u_{k+j}) = p(\bar{\gamma})}$$

where  $u_i, u_{k+j} \in H^4$  are the first Pontryagin classes of the canonical rank-2 vector bundles over  $O(2n+2)/(O(2)^{2k} \times O(1) \times O(2)^{2n-2k} \times O(1))$ , and  $\gamma, \bar{\gamma}$  are the pull-backs to  $O(2n+2)/(O(2)^{2k} \times O(1) \times O(2)^{2n-2k} \times O(1))$  of the canonical and complementary bundles over the Grassmannian  $G_{2k+1}(\mathbb{R}^{2n+2})$ . Moreover, the Poincaré series are  $P_{G_{2k+1}(\mathbb{R}^{2n+2})}(t) \prod_{i=1}^k \frac{1-t^{4i}}{1-t^4} \prod_{j=1}^{n-k} \frac{1-t^{4j}}{1-t^4}$ .

On the other hand, we can consider the following different fibration on the same manifold  $O(2n+2)/(O(2)^{2k} \times O(1) \times O(2)^{2n-2k} \times O(1))$ :

$$\frac{O(2n+1)}{O(2)^{2n} \times O(1)} \hookrightarrow \frac{O(2n+2)}{O(2)^{2k} \times O(1) \times O(2)^{2n-2k} \times O(1)} \rightarrow \frac{O(2n+2)}{O(2n+1) \times O(1)}$$

where the fibre is the complete even-rank flag manifold  $Fl(\mathbb{R}^{2n+1})$  and the base is  $\mathbb{R}P^{2n+1}$ . Applying the Proposition 2.2.2 to this complete even-rank flag bundle, we get:

$$H^*\left(\frac{O(2n+2)}{O(2)^{2k} \times O(1) \times O(2)^{2n-2k} \times O(1)}, \mathbb{Q}\right) \cong \frac{H^*(\mathbb{R}P^{2n+1}, \mathbb{Q})[u_1, \dots, u_n]}{\prod_{i=1}^n (1+u_i) = 1} \cong \frac{\mathbb{Q}[r, u_1, \dots, u_n]}{r^2 = 0, \prod_{i=1}^n (1+u_i) = 1}$$

where  $r \in H^{2n+1}$  is the unit orientation form on  $\mathbb{R}P^{2n+1}$ . Moreover, the Poincaré series are  $(1+t^{2n+1}) \prod_{i=1}^n \frac{1-t^{4i}}{1-t^4}$ .

Comparing the two ways of computing Poincaré series, we get

$$P_{G_{2k+1}(\mathbb{R}^{2n+2})}(t) = (1+t^{2n+1}) \frac{(1-t^4) \dots (1-t^{4n})}{(1-t^4) \dots (1-t^{4k})(1-t^4) \dots (1-t^{4n-4k})} = (1+t^{2n+1}) P_{G_{2k}(\mathbb{R}^{2n})}(t)$$

Comparing the two ways of computing cohomology rings, we get:

$$\frac{H^*(G_{2k+1}(\mathbb{R}^{2n+2}), \mathbb{Q})[u_1, \dots, u_k; u_{k+1}, \dots, u_n]}{\prod_{i=1}^k (1+u_i) = p(\gamma), \prod_{j=1}^{n-k} (1+u_{k+j}) = p(\bar{\gamma})} \cong \frac{\mathbb{Q}[r, u_1, \dots, u_n]}{r^2 = 0, \prod_{i=1}^n (1+u_i) = 1}$$

which means the odd-degree element  $r$  comes from  $H^*(G_{2k+1}(\mathbb{R}^{2n+2}), \mathbb{Q})$ , and so do the relations  $r^2 = 0, p(\gamma)p(\bar{\gamma}) = 1$ . Similarly, we can compare two different fibrations on the same homogeneous space  $SO(2n+2)/(SO(2)^{2k} \times SO(1) \times SO(2)^{2n-2k} \times SO(1))$ . This is essentially Takeuchi's result:

**Proposition 2.3.1** (Cohomology rings of odd dimensional Grassmannians, [Ta62]). *The cohomology rings of  $G_{2k+1}(\mathbb{R}^{2n+2}), \tilde{G}_{2k+1}(\mathbb{R}^{2n+2})$  are generated by Pontryagin classes and a new type of odd-degree classes  $r, \tilde{r} \in H^{2n+1}$ :*

$$\begin{aligned} H^*(G_{2k+1}(\mathbb{R}^{2n+2}), \mathbb{Q}) &= \frac{\mathbb{Q}[p_1, p_2, \dots, p_k; \bar{p}_1, \bar{p}_2, \dots, \bar{p}_{n-k}; r]}{p\bar{p} = 1, r^2 = 0} \\ H^*(\tilde{G}_{2k+1}(\mathbb{R}^{2n+2}), \mathbb{Q}) &= \frac{\mathbb{Q}[p_1, p_2, \dots, p_k; \bar{p}_1, \bar{p}_2, \dots, \bar{p}_{n-k}; \tilde{r}]}{p\bar{p} = 1, \tilde{r}^2 = 0} \end{aligned}$$

Moreover, the same argument also gives a bundle version:

- (1) If  $V \rightarrow M$  is a real vector bundle of rank  $2n+2$ , let  $\mathbb{R}P(V)$  be its real projectivization, then we get the isomorphism of cohomology rings

$$H^*(G_{2k+1}(V^{2n+2}), \mathbb{Q}) \cong \frac{H^*(\mathbb{R}P(V^{2n+2}), \mathbb{Q})[p_1, p_2, \dots, p_k; \bar{p}_1, \bar{p}_2, \dots, \bar{p}_{n-k}]}{p\bar{p} = p(V)}$$

- (2) If  $V \rightarrow M$  is an oriented vector bundle of rank  $2n + 2$ , let  $S(V)$  be its sphere bundle, then then we get the isomorphism of cohomology rings

$$H^*(\tilde{G}_{2k+1}(V^{2n+2}), \mathbb{Q}) \cong \frac{H^*(S(V^{2n+2}), \mathbb{Q})[p_1, p_2, \dots, p_k; \bar{p}_1, \bar{p}_2, \dots, \bar{p}_{n-k}]}{p\bar{p} = p(V)}$$

*Remark 2.3.1.* To relate the cohomology of  $G_{2k+1}(V^{2n+2})$  or  $\tilde{G}_{2k+1}(V^{2n+2})$  with the cohomology of  $M$ , because of the above isomorphisms, we need to relate the cohomology of  $\mathbb{R}P(V^{2n+2})$  or  $S(V^{2n+2})$  with the cohomology of  $M$  by applying the Gysin sequence to the bundle  $\mathbb{R}P(V) \rightarrow M$  or  $S(V) \rightarrow M$ . However, the Gysin sequence only works for group structure not for ring structure.

**2.3.3. Associated equivariant flag bundle.** Suppose  $V \rightarrow B$  is an equivariant vector bundle acted by a torus  $T$ , then the associated flag bundles are also  $T$ -equivariant. Besides the ordinary cohomology of the associated flag bundles, we can also consider their  $T$ -equivariant cohomology. Without loss of generality, we will give equivariant cohomology ring of associated equivariant complete (even-rank) flag bundle and Grassmannian bundle. Here, we use the Borel model of equivariant cohomology  $H_T^*(M) = H^*(M_T)$ , where the Borel construction  $M_T = ET \times_T M$  is a bundle over  $BT$  with fibre  $M$ .

We follow Mare's observation (see [Ma]): the Borel construction of associated flag bundle is the associated flag bundle of the Borel construction. For example,  $(G_k(V))_T = ET \times_T G_k(V) \cong G_k(ET \times_T V)$ . Then we can apply the Theorem 2.2.1, and obtain similar expressions with ordinary characteristic classes replaced by equivariant characteristic classes:

- (1) If  $V^n$  is a  $T$ -equivariant rank- $n$  complex vector bundle, then we get the equivariant cohomology ring

$$H_T^*(G_k(V^n), \mathbb{Z}) \cong \frac{H_T^*(B, \mathbb{Z})[c_1^T, c_2^T, \dots, c_k^T; \bar{c}_1^T, \bar{c}_2^T, \dots, \bar{c}_{n-k}^T]}{c^T \bar{c}^T = c^T(V^n)}$$

- (2) If  $V$  is a  $T$ -equivariant real vector bundle of rank  $2n$  or  $2n + 1$ , denoted as  $V^{2n}$  or  $V^{2n+1}$  respectively, then we get the equivariant cohomology rings

$$\begin{aligned} H_T^*(G_{2k}(V^{2n}), \mathbb{Q}) &\cong \frac{H_T^*(B, \mathbb{Q})[p_1^T, p_2^T, \dots, p_k^T; \bar{p}_1^T, \bar{p}_2^T, \dots, \bar{p}_{n-k}^T]}{p^T \bar{p}^T = p^T(V^{2n})} \\ H_T^*(G_{2k}(V^{2n+1}), \mathbb{Q}) &\cong H_T^*(G_{2k+1}(V^{2n+1}), \mathbb{Q}) \cong \frac{H_T^*(B, \mathbb{Q})[p_1^T, p_2^T, \dots, p_k^T; \bar{p}_1^T, \bar{p}_2^T, \dots, \bar{p}_{n-k}^T]}{p^T \bar{p}^T = p^T(V^{2n+1})} \end{aligned}$$

- (3) If  $V$  is a  $T$ -equivariant oriented vector bundle of rank  $2n$  or  $2n + 1$ , denoted as  $V^{2n}$  or  $V^{2n+1}$  respectively, then we get the equivariant cohomology rings

$$\begin{aligned} H_T^*(\tilde{G}_{2k}(V^{2n}), \mathbb{Q}) &\cong \frac{H_T^*(B, \mathbb{Q})[p_1^T, p_2^T, \dots, p_k^T; \bar{p}_1^T, \bar{p}_2^T, \dots, \bar{p}_{n-k}^T; e^T, \bar{e}^T]}{p^T \bar{p}^T = p^T(V^{2n}), (e^T)^2 = p_k^T, (\bar{e}^T)^2 = \bar{p}_{n-k}^T, e^T \bar{e}^T = e^T(V^{2n})} \\ H_T^*(\tilde{G}_{2k}(V^{2n+1}), \mathbb{Q}) &\cong \frac{H_T^*(B, \mathbb{Q})[p_1^T, p_2^T, \dots, p_k^T; \bar{p}_1^T, \bar{p}_2^T, \dots, \bar{p}_{n-k}^T; e^T]}{p^T \bar{p}^T = p^T(V^{2n+1}), (e^T)^2 = p_k^T} \\ H_T^*(\tilde{G}_{2k+1}(V^{2n+1}), \mathbb{Q}) &\cong \frac{H_T^*(B, \mathbb{Q})[p_1^T, p_2^T, \dots, p_k^T; \bar{p}_1^T, \bar{p}_2^T, \dots, \bar{p}_{n-k}^T; \bar{e}^T]}{p^T \bar{p}^T = p^T(V^{2n+1}), (\bar{e}^T)^2 = \bar{p}_{n-k}^T} \end{aligned}$$

Similarly, we can get the equivariant cohomology ring of associated complete (even-rank) flag bundle by replacing the ordinary cohomology classes in Proposition 2.2.2 with equivariant cohomology classes. Same idea applies to the cohomology ring of associated partial (even-rank) flag bundles.

As a special case, if we take the base  $B$  to be a point, and note  $H_{T^n}^*(pt; R) = R[\alpha_1, \dots, \alpha_n]$  where  $\alpha_1, \dots, \alpha_n$  are the standard integral basis of the dual Lie algebra  $\mathfrak{t}^*$ , then we can get the equivariant cohomology ring of complete (even-rank) flag manifolds:

- (1) If  $V = \mathbb{C}^n$ , then we get the equivariant cohomology ring for the complete complex flag manifold

$$H_{T^n}^*(Fl(\mathbb{C}^n), \mathbb{Z}) = \frac{\mathbb{Z}[\alpha_1, \alpha_2, \dots, \alpha_n][x_1, x_2, \dots, x_n]}{\prod_{i=1}^n (1 + x_i) = \prod_{i=1}^n (1 + \alpha_i)}$$

where  $x_i \in H_{T^n}^2$  is the equivariant first Chern class of the  $i$ -th canonical complex line bundle over  $Fl(\mathbb{C}^n)$ .

- (2) If  $V = \mathbb{R}^{2n}$  or  $\mathbb{R}^{2n+1}$ , then we get the equivariant cohomology rings for the complete even-rank flag manifolds

$$\begin{aligned} H_{T^n}^*(Fl(\mathbb{R}^{2n}), \mathbb{Q}) &= \frac{\mathbb{Q}[\alpha_1, \alpha_2, \dots, \alpha_n][u_1, u_2, \dots, u_n]}{\prod_{i=1}^n (1 + u_i) = \prod_{i=1}^n (1 + \alpha_i^2)} \\ H_{T^n}^*(Fl(\mathbb{R}^{2n+1}), \mathbb{Q}) &= \frac{\mathbb{Q}[\alpha_1, \alpha_2, \dots, \alpha_n][u_1, u_2, \dots, u_n]}{\prod_{i=1}^n (1 + u_i) = \prod_{i=1}^n (1 + \alpha_i^2)} \end{aligned}$$

where  $u_i \in H_{T^n}^4$  is the equivariant first Pontryagin class of the  $i$ -th canonical rank-2 bundle over  $Fl(V)$ .

- (3) If  $V = \mathbb{R}^{2n}$  or  $\mathbb{R}^{2n+1}$  with the natural orientation, then we get the equivariant cohomology rings for the complete even-rank oriented flag manifolds

$$\begin{aligned} H_{T^n}^*(\tilde{Fl}(\mathbb{R}^{2n}), \mathbb{Q}) &= \frac{\mathbb{Q}[\alpha_1, \alpha_2, \dots, \alpha_n][u_1, u_2, \dots, u_n; e_1, e_2, \dots, e_n]}{\prod_{i=1}^n (1 + u_i) = \prod_{i=1}^n (1 + \alpha_i^2), e_i^2 = u_i, \prod_{i=1}^n e_i = \prod_{i=1}^n \alpha_i} \\ &= \frac{\mathbb{Q}[\alpha_1, \alpha_2, \dots, \alpha_n][e_1, e_2, \dots, e_n]}{\prod_{i=1}^n (1 + e_i^2) = \prod_{i=1}^n (1 + \alpha_i^2), \prod_{i=1}^n e_i = \prod_{i=1}^n \alpha_i} \\ H_{T^n}^*(\tilde{Fl}(\mathbb{R}^{2n+1}), \mathbb{Q}) &= \frac{\mathbb{Q}[\alpha_1, \alpha_2, \dots, \alpha_n][u_1, u_2, \dots, u_n; e_1, e_2, \dots, e_n]}{\prod_{i=1}^n (1 + u_i) = \prod_{i=1}^n (1 + \alpha_i^2), e_i^2 = u_i} \\ &= \frac{\mathbb{Q}[\alpha_1, \alpha_2, \dots, \alpha_n][e_1, e_2, \dots, e_n]}{\prod_{i=1}^n (1 + e_i^2) = \prod_{i=1}^n (1 + \alpha_i^2)} \end{aligned}$$

where  $e_i \in H_{T^n}^2$  is the equivariant Euler class of the  $i$ -th canonical rank-2 bundle over  $\tilde{Fl}(V)$ .

**2.3.4. Bott tower of flag manifolds.** In the proof of Theorem 2.2.1, we used the fact that associated complete (even-rank) flag bundle is obtained through a sequence of projectivizations or rank-2 Grassmannianizations. We can consider general Bott tower of associated flag bundles.

$$\begin{array}{ccccccc} & V_m & & \cdots & & V_3 & & V_2 & & V_1 \\ & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ E_m & \longrightarrow & E_{m-1} & \longrightarrow & \cdots & \longrightarrow & E_2 & \longrightarrow & E_1 & \longrightarrow & E_0 = B \end{array}$$

where each  $E_i$  is obtained successively as an associated partial (even-rank) flag bundle of the vector bundle  $V_i \rightarrow E_{i-1}$ .

In the simplest case, we start with  $E_0$  as a point and construct  $E_i \rightarrow E_{i-1}$  as the associated projectivization of a complex rank-2 vector bundle  $V_i \rightarrow E_{i-1}$ . This is the Bott tower of  $\mathbb{C}P^1$ , starting at a point. For each projectivization, we get

$$H^*(E_i, \mathbb{Z}) \cong \frac{H^*(E_{i-1}, \mathbb{Z})[x_i]}{x_i^2 - c_1(V_i)x_i + c_2(V_i) = 0}$$

Therefore,

$$H^*(E_m, \mathbb{Z}) \cong \frac{\mathbb{Z}[x_1, x_2, \dots, x_m]}{x_i^2 - c_1(V_i)x_i + c_2(V_i) = 0, \quad 1 \leq i \leq m}$$

In the general case, we can still use the Leray-Hirsch Theorem 2.1.2 and the Remark 2.2.2 successively to get the cohomology ring of  $E_i$  as a ring extension of the cohomology of  $E_{i-1}$ . Furthermore, if the vector bundle  $V_1 \rightarrow E_0$  admits a  $T$ -action that can be lifted successively to  $V_i \rightarrow E_{i-1}$ , then we can also compute the

equivariant cohomology successively using the previous discussion of associated equivariant partial (even-rank) flag bundle.

### 3. FINAL DISCUSSION

The author is gratefully indebted to Jeffrey Carlson for pointing out that the calculation of cohomology ring of associated complex, real or oriented flag bundle can be strengthened to associated bundle with fibre being generalized flag manifolds using a spectral sequence argument of Larry Smith.

**Theorem 3.0.1** (Cohomology ring of pullback bundle, [Sm67]). *Let  $F \hookrightarrow E_0 \rightarrow B_0$  be a fibre bundle and suppose we are given a pull-back diagram*

$$\begin{array}{ccc} E_1 & \longrightarrow & E_0 \\ \downarrow & & \downarrow \\ B_1 & \longrightarrow & B_0 \end{array}$$

*If the Serre spectral sequence of  $F \hookrightarrow E_0 \rightarrow B_0$  collapses at  $E_2$  page, then*

- (a) *The Serre spectral sequence of  $F \hookrightarrow E_1 \rightarrow B_1$  also collapses at  $E_2$  page*
- (b) *The cohomology rings fit into a push-out diagram:*

$$\begin{array}{ccc} H^*(E_1) & \longleftarrow & H^*(E_0) \\ \uparrow & & \uparrow \\ H^*(B_1) & \longleftarrow & H^*(B_0) \end{array}$$

*Or equivalently,  $H^*(E_1) \cong H^*(B_1) \otimes_{H^*(B_0)} H^*(E_0)$  as a ring.*

*Remark 3.0.2.* The original proof of Smith uses Serre spectral sequences and Eilenberg-Moore spectral sequences. Carlson has a short proof using only Serre spectral sequences in his thesis [Carl15].

**Corollary 3.0.1** (Cohomology ring of associated generalized flag bundle, [Carl]). *Let  $G$  be a compact connected Lie group and  $H$  be a closed subgroup of  $G$  with the same rank. For any principal  $G$ -bundle  $P \rightarrow B$ , the fibre bundle  $P/H \rightarrow B$  has the generalized flag manifold  $G/H$  as fibres with cohomology ring given as*

$$H^*(P/H) \cong H^*(B) \otimes_{H^*(BG)} H^*(BH)$$

*Moreover, if the principal  $G$ -bundle  $P \rightarrow B$  admits an extra action by a compact Lie group  $K$  that commutes with the  $G$ -action, then the  $K$ -equivariant cohomology of  $P/H$  can be given as*

$$H_K^*(P/H) \cong H_K^*(B) \otimes_{H^*(BG)} H^*(BH)$$

*Proof.* The principal  $G$ -bundle  $P \rightarrow B$  is a pull-back of the universal principal  $G$ -bundle  $EG \rightarrow BG$  via a classifying map  $B \rightarrow BG$ , i.e.

$$\begin{array}{ccc} P & \longrightarrow & EG \\ \downarrow & & \downarrow \\ B & \longrightarrow & BG \end{array}$$

Take quotients by  $H$  on  $P$  and  $EG$ , and note that  $(EG)/H = (EH)/H = BH$ , then we get a pull-back diagram

$$\begin{array}{ccc} P/H & \longrightarrow & BH \\ \downarrow & & \downarrow \\ B & \longrightarrow & BG \end{array}$$

Since both bundles  $P/H \rightarrow B$  and  $BH \rightarrow BG$  have the same fibre  $G/H$  whose cohomology concentrates at even degrees in appropriate coefficients, the Serre spectral sequences of these two bundles collapse at  $E_2$  page. Then the Smith's Theorem 3.0.1 gives the cohomology ring of the associated generalized flag bundle  $P/H$ . The  $K$ -equivariant version is obtained by considering the  $G$ -principal Borel construction  $EK \times_K P \rightarrow EK \times_K B$ .  $\square$

As a special case of Smith's general theorem, our survey gives slightly more details on the generators and relations of the cohomology  $H^*(P/H)$  as a ring extension of the cohomology  $H^*(B)$  for  $G$  being  $U(n)$ ,  $O(n)$ ,  $SO(n)$ . In this case, the method of splitting principle is perhaps more user-friendly than the language of spectral sequences.

## REFERENCES

- [Bo53a] A. Borel, Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts. *Ann. of Math.* (2) 57, (1953). 115-207.
- [Bo53b] A. Borel, La cohomologie mod 2 de certains espaces homogènes, *Comment. Math. Helv.* 27, (1953). 165-197.
- [BT82] R. Bott and L. Tu, *Differential forms in algebraic topology*, Graduate Texts in Mathematics, 82. Springer-Verlag, New York-Berlin, 1982.
- [Ca78] J. Carrell, Chern classes of the Grassmannians and Schubert calculus. *Topology* 17 (1978), no. 2, 177-182.
- [Car50] H. Cartan, La transgression dans un groupe de Lie et dans un espace fibré principal, *Colloque de topologie (espaces fibrés)*, Bruxelles, 1950, pp. 57-71.
- [Carl] J. Carlson, Personal Communication.
- [Carl15] J. Carlson, On the equivariant cohomology of homogeneous spaces, Ph.D. Thesis, Tufts University. 2015. 345 pp.
- [CK] L. Casian and Y. Kodama, On the cohomology of real Grassmann manifolds, preprint, <https://arxiv.org/pdf/1309.5520v1.pdf>
- [Ha02] A. Hatcher, *Algebraic topology*, Cambridge University Press, Cambridge, 2002. xii+544 pp.
- [He] C. He, GKM theory, characteristic classes and the equivariant cohomology ring of real Grassmannian, preprint, <https://arxiv.org/pdf/1609.06243v2.pdf>
- [Le49] J. Leray, Détermination dans les cas non exceptionnels de l'anneau de cohomologie de l'espace homogène quotient d'un groupe de Lie compact par un sous-groupe de même rang, *C. R. Acad. Sci. Paris*, 228, (1949). 1902-1904.
- [Ma] A.-L. Mare, Equivariant cohomology of complex flag manifolds, preprint, <http://uregina.ca/~mareal/equivcohfl.pdf>
- [Ja89] J. Jaworowski, An additive basis for the cohomology of real Grassmannians. *Algebraic topology Poznan 1989*, 231-234, *Lecture Notes in Math.*, 1474, Springer, Berlin, 1991.
- [MS74] J. Milnor and J. Stasheff, *Characteristic classes*. Annals of Mathematics Studies, No. 76. Princeton University Press, Princeton, N. J.; University of Tokyo Press, Tokyo, 1974.
- [Sm67] L. Smith, Homological algebra and the Eilenberg-Moore spectral sequence, *Trans. Amer. Math. Soc.* 129 1967 58-93.
- [Ta62] M. Takeuchi, On Pontryagin classes of compact symmetric spaces. *J. Fac. Sci. Univ. Tokyo Sect. I*, 9 (1962) 313-328.

DEPARTMENT OF MATHEMATICS, NORTHEASTERN UNIVERSITY, BOSTON, USA

E-mail address: [he.chen@husky.neu.edu](mailto:he.chen@husky.neu.edu)